

## Biodegradation of Tricalcium Phosphate by Phosphate Solubilizing Bacteria

<sup>1</sup>V. Madhan Chakkaravarthy, <sup>3</sup>R. Arunachalam, <sup>2</sup>S. Vincent, <sup>3</sup>K. Paulkumar and <sup>3</sup>G. Annadurai

<sup>1</sup>Zoological Survey of India, Andaman and Nicobar Regional Centre,  
National Coral Reef Research Institute, Port Blair-744 102, A and N Island, India

<sup>2</sup>Loyola Institute of Frontier Energy, Loyola College, Chennai-600 034, India

<sup>3</sup>Environmental Nanotechnology Division, Sri Paramakalyani Centre for Environmental Sciences,  
Manonmaniam Sundaranar University, Alwarkurichi-627 412, Tamilnadu, India

**Abstract:** The present study was aimed at to develop a Phosphate Solubilizing Bacteria's (PSB) inoculum for improving Phosphate (P) uptake on phosphate sludge. Twelve PSB were isolated from three types of rhizosphere soil from Chennai, India were studied for phosphate solubilization on Pikovskaya's medium. Isolates were subjected and studied for Solubilization Index (SI) for selecting the high efficient strain. In results, the PB08 strain exhibited high SI 4.80 at sixth day incubation and it was grown *in vitro* for ten days in Pikovskaya's broth containing 1% tricalcium phosphate (TCP) and 1% Phosphate Sludge (PS). Variation occurs in phosphate solubilization of two media which contain two P source inoculated by the same strain (PB08). Unlike SI and pH drop, fluctuations in phosphate solubilization were observed during 10 days and gradual increase in phosphate solubilization was noted. Values of solubilized tricalcium phosphate ranged from 9.6 to 136 ppm. The results emphasize that the PB08 may be used for inoculum production and their inoculation effect on plant growth can be studied *in vivo*.

**Key words:** Biodegradation, phosphate solubilizing bacteria, tricalcium phosphate, phosphate sludge, solubilization index

### INTRODUCTION

Phosphorus (P) is the second most important plant nutrient after nitrogen (Donahue *et al.*, 1990). However, many soils throughout the world are P deficient because the free phosphorus concentration even in fertile soils is generally not higher than 10  $\mu\text{M}$  even at 6.5 pH where it is most soluble. To circumvent the P deficiency in soils, P fertilizers are applied. However, after application, a considerable amount of P is rapidly transformed into less available forms by forming a complex with Al or Fe in acid soils or Ca in calcareous soils (Lindsay *et al.*, 1989) before plant roots had a chance to absorb it (Vikram, 2007).

Numerous microorganisms, especially those associated with roots have the ability to increase plant growth and productivity (Chang *et al.*, 1986; Kloepper *et al.*, 1988). Phosphate-solubilizing microbes play fundamental roles in biogeochemical phosphorus cycling in natural and agricultural ecosystems. Phosphate-solubilizing microbes can transform the insoluble phosphorus to soluble forms  $\text{HPO}_4^{2-}$  and  $\text{H}_2\text{PO}_4^-$  by acidification, chelation, exchange reactions and polymeric substances formation (Chang and Yang,

2009; Delvasto *et al.*, 2006). Therefore, the use of phosphate-solubilizing microbes in agricultural practice would not only offset the high cost of manufacturing phosphatic fertilizers but would also mobilize insoluble phosphorus in the fertilizers and soils to which they are applied (Chang and Yang, 2009; Rodriguez and Fraga, 1999). Application of the phosphate-solubilizing microbes *Agrobacterium*, *Bacillus*, *Enterobacter*, *Pseudomonas*, *Aspergillus*, *Trichoderma* and *Glomus* around the roots of plants, in soils and in fertilizers has been shown to release soluble phosphorus, promote plant growth and protect plants from pathogen infection (Chang and Yang, 2009; Rodriguez and Fraga, 1999; Rudresh *et al.*, 2005; Zayed and Abdel-Motaal, 2005a, b; Biswas and Narayanasamy, 2006; Ouahmane *et al.*, 2007). Therefore, removal of phosphorus is prerequisite for restoration of water quality (Van and Boers, 1994).

Bacteria of the genus *Bacillus* are ubiquitous and common soil microorganisms that play an important role in silicate biodegradation during the process of rock disintegration (Vikram and Hamzehzarghani, 2008; Girgis *et al.*, 2008; Han *et al.*, 2006; Liu *et al.*, 2006). The results of such activity involve both geochemical and

structural changes in rocks and silicate and the most powerful phosphate solubilizers (Rodriguez and Fraga, 1999). The metabolic diversity of *Bacillus* sp. in particular together with its low reported incidence of pathogenicity, has led to the fact that many representatives of this group are being used in a wide range of applications. Due to its ability to produce a range of enzymes, solubilization pounded nutrients and degrade organic wastes along with the N-fixing ability of some strains, *Bacillus* sp seemed to be a good candidate for biofertilizers application in agriculture (Girgis *et al.*, 2008). Inoculation with bacteria, which can improve P and K availability in soils by producing organic acids and other chemicals, stimulated growth and mineral uptake of plants (Girgis *et al.*, 2008; Garcia *et al.*, 2004). Hence, the present study has been designed to develop a PSB inoculum for improving P uptake on phosphate sludge.

## MATERIALS AND METHODS

**Isolation and cultivation of PSB:** The plants of groundnut, sorghum and cotton were uprooted from field soils which were obtained from Chennai, India and the soil loosely adhered to roots was removed by mechanical shaking. The investigation was performed during November 2008 to February 2009. Each soil sample (0.2 g) was placed in an Erlenmeyer flask (250 mL) containing 100 mL of sterilized Pikovskaya's liquid medium. The flasks were incubated for 7 days on an orbital shaker at 200 rpm at room temperature (23±4°C). Pikovskaya's Broth (PB) consisted of the following: Ca<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub> 2.5 g, glucose 13 g, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> 0.5 g, NaCl 0.2 g, MgSO<sub>4</sub>·7H<sub>2</sub>O 0.1 g, KCl 0.2 g, yeast extract 0.5 g, MnSO<sub>4</sub> trace, FeSO<sub>4</sub>·7H<sub>2</sub>O trace and dissolved in 1000 mL distilled water. The pH adjusted to 7.2 insoluble tricalcium phosphates was used as a growth limiting factor as a result PSMs could continue growth. After 7 days of incubation in liquid medium, the culture was serially diluted and streaked on Pikovskaya's solid medium (PA) containing 1.5% agar. Twelve bacterial species were isolated through successive cultivations and microorganism isolated from the rhizospheric soil of groundnut showed the best phosphate solubilization on agar medium.

**Solubilization Index (SI):** The 0.1 mL of each PSB culture preserved in sterile distilled water was placed on Pikovskaya's agar plate and incubated for seven days. Solubilization index was measured using the following formula (Premono *et al.*, 1996). The PB08 showed the best SI on agar plate and it was temporarily identified as *Bacillus* sp. by the analysis of gram reaction, sugar fermentation, starch and casein hydrolysis (data not shown).

$$SI = \frac{\text{Colony diameter} + \text{Halozone diameter}}{\text{Colony diameter}}$$

**pH change:** In medium acidification of culture was visualized by spot inoculation on Pikovskaya's medium containing bromothymol blue and bromocresol purple indicator. A 1 mL of three day old culture of PB08 (containing about 1×10<sup>3</sup> cfu) was added to two sterile 250 mL Pikovskaya's broth containing 1% tricalcium phosphate and 1% Phosphate Sludge (PS) and kept on shaker for ten days. Sterile uninoculated medium served as control. Initial pH and changes in pH were noted each day by digital pH meter.

**Determination of phosphate solubilization:** A 10 mL sample of each culture was centrifuged for 15 min at 1500 rpm. Phosphate in solution was extracted with Ammonium bicarbonate diethylene triamine penta acetic acid (Soltanpour and Workman, 1979). Supernatant was decanted and 5 mL of supernatant was added to 20 mL of AB-DTPA extracting solution. The mixture was shaken on a reciprocating shaker for 15 min at 180 cpm in open flasks and the extract was stored in plastic bottle. One milliliter of chloromolybdic acid was added and diluted with 25-30 mL of distilled water, then 0.25 mL of chlorostannous acid was added and the volume was made up to 50 mL with distilled water. The solution was shaken thoroughly and the blue colour developed was read at 660 nm using spectrophotometer and the concentration of available phosphate was calculated.

## RESULTS

Solubilization index based on colony diameter and halozone for each isolate is presented in Fig. 1a and b, results shows that PB08 was most efficient phosphate solubilizer on phosphate agar plates with SI = 4.80 at sixth day incubation. Studies on agar plates revealed that phosphate solubilizing bacteria formed clear zones by solubilizing suspended tricalcium phosphate (data not shown). Measurements of SI ranged from 2-4.8. Generally, halozone increased with increase in colony diameter. Fluctuations in solubilization index were observed during the seven day observation period. In most of the cases it gradually increased, while in few cases (PB01, PB03, PB04 and PB011) increased initially and later decreased. Most of PSB strains lost their ability to form halozones on phosphate agar medium on repeated subculturing. Microbial phosphate solubilization on plant roots has been by agar contact method with a pH indicator (bromocresol purple) has also been developed for the detection of zone of acidification in the rhizosphere and




Fig. 1: (a) pH of both the medium (PB+TCP and PB+PS) incubated for ten days. (b) Variation in phosphate solubilization of two media incubated for ten days

could be a valuable tool for studying phosphate solubilizing microorganisms. In the present study the diffusion of acid into the medium was visualized by the presence of bromocresol purple. It was also reflected in the diameter of zone of calcium phosphate solubilization.

The PB08 showed the best SI and it was identified as the genus *Bacillus* PB08 lowered the pH in both the medium (PB+TCP and PB+PS) as compared to uninoculated sterile PB control incubated for ten days under conditions as inoculated (Fig. 1a). Fluctuations in pH drop were also noted. pH studies showed a drop of pH from 7.04 (control) to 3.0 (TCP) at the 3rd day and 3.38 (PS) at 9th day (Fig. 1a).

The phosphate solubilization the bacterial isolate PB08 was solubilized phosphate in Pikovskaya's medium which contains TCP and PS. Variation occurs in phosphate solubilization of two media which contain two P source inoculated by the same strain (PB08) (Fig. 1b). Unlike SI and pH drop, fluctuations in phosphate solubilization were observed during 10 days and gradual increase in phosphate solubilization was noted. Values of solubilized tricalcium phosphate ranged from 9.6 to 133.6 ppm for TCP and 12.6 to 136 ppm for PS. In both the cases increased phosphate solubilization was noted at 10th day (Fig. 1b).

## DISCUSSION

Girgis *et al.* (2008) reports showed that the increasing of total acidity percentage was responsible in decreasing the pH. Lowest total acidity percentage was recorded in a range of 0.41 to 0.44 and it reveals the decrease in the pH values in MA-m or MA-f culture media with increase in their total acidity percentages was not the only direct reason for the release of soluble K. However, no relationships were observed between the final pH and the percentages of total acidity in their culture media and the amount of soluble K. On the other side, the decrease in pH with the increase of total acidity in PVK-tcp or PVK-rp (6.68%) culture media may explain why higher concentration of released P was detected. Furthermore, linkage was observed between the final pH and the total acidity of the culture media and the amount of soluble P. In this concern, Groudeva and Groudev (1987) noted that the bacterial action on silicate and aluminosilicate is connected with the formation of mucilaginous capsules consisting of EPS as well as the production of different metabolites such as organic and amino acids. They did not exclude that the bacterial action may also be resulting of an enzymatic nature. Styriakova *et al.* (2004) reported that the activity of silicate dissolving bacteria played a stimulation role in the release of Si, Fe<sup>3+</sup> and K<sup>+</sup> from

feldspar and Fe<sup>3+</sup> oxyhydroxides. The binding of silicate to the bacterial surfaces can thus be described as an outer sphere complex formation as it occurs through electrostatic interaction. The present investigation shows that PB08 was most efficient phosphate solubilizer on phosphate agar plates with SI = 4.80 at sixth day incubation (Fig. 1). Studies on agar plates revealed that phosphate solubilizing bacteria formed clear zones by solubilizing suspended tricalcium phosphate. De-Souza *et al.* (2000) results showed that the five of the 88 phosphate solubilizing isolates showed higher phosphatase activity (8.68-24.78 µmol/mL/day) with culture P63 showing the highest activity. In present study twelve PSB were isolated from three types of rhizosphere soil from Chennai, India were studied for phosphate solubilization on Pikovskaya's medium. Isolates were subjected and studied for Solubilization Index (SI) for selecting the high efficient strain. Variation occurs in phosphate solubilization of two media which contain two P source inoculated by the same strain (PB08) (Fig. 1b). Unlike SI and pH drop, fluctuations in phosphate solubilization were observed during 10 days and gradual increase in phosphate solubilization was noted. Values of solubilized tricalcium phosphate ranged from 9.6 to 136 ppm (Fig. 1b).

Seshadri *et al.* (2002) reported the solubilizing index of *Bacillus* sp., was 184.19±6.43 at 7.83-8.21 pH. Present study showed the initial solubilization index of *Bacillus* sp., was 4.80 at 6th day at 3-7.04 pH in PB+TCP and 3.8-7.04 in PB+PS. At above the pH PB+TCP *Bacillus* sp., showed the solubilization 9.6-133.6 and PB+PS showed 12.6-136 (ppm). The pH used for the present investigation is lower than the report of Seshadri *et al.* (2002), it suggest that in acidic condition also suitable for P solubilization. The maximum solubilization was observed in PB+TCP acidic medium that was 133.6 and PB+PS was 136 ppm at 10th day. These results clearly suggest the *Bacillus* sp., culture medium in alkaline condition most suitable for P solubilization than medium in acidic nature.

Lebo (1990) reported that PSB were not restricted to a few genera as it mainly depends on the regions. In this study the strains were mainly *Bacillus*, in addition few strains of *Vibrio* and *Klebsiella* were also encountered, whereas in oceanic regions *Pseudomonas* was the dominant group and in the Tokyo Bay it was *Flavobacterium* and *Acinetobacter*. These results highly support to the present study by the similar organism *Bacillus* sp., was used for decomposing tribase calcium phosphate. *Pseudomonas* and *Bacillus* sp., were found to have more phosphate solubilizing capacity than the other genera (*Vibrio*, *Alcaligenes* and *Corynebacterium*) (Seshadri *et al.*, 2002). Venkateswaran and Natarajan

(1983) while studying the Porto Novo waters reported *Pseudomonas* sp. and *Bacillus* sp., as dominant inorganic phosphorus compounds solubilizing microbes. These results reveal the *Bacillus* sp., found to be an effective microbe for phosphate solubilizing activities.

## CONCLUSION

It is concluded from the present study that the fluctuations occur in SI and acid production (pH drop) but not in phosphate solubilization. Present study also showed that PB08 is the most efficient strain on the basis of its phosphate solubilizing activity. Further research should be continued with such efficient strain. This may be used for inoculum production and its inoculation effect on the plant growth may be studied *in vivo*.

## REFERENCES

- Biswas, D.R. and G. Narayanasamy, 2006. Rock phosphate enriched compost: An approach to improve low-grade Indian rock phosphate. *Bioresour. Technol.*, 97: 2243-2251.
- Chang, Y.C., Y.C. Chang, R. Baker, O. Kleifeld and I. Chet, 1986. Increased growth of plants in the presence of the biological control agents *Trichoderma harzianum*. *Plant Dis.*, 70: 145-148.
- Chang, C.H. and S.S. Yang, 2009. Thermo-tolerant phosphate-solubilizing microbes for multi-functional biofertilizer preparation. *Bioresour. Technol.*, 100: 1648-1658.
- De-Souza, M.J.B.D., S. Nair and D. Chandramohan, 2000. Phosphate solubilizing bacteria around Indian peninsula. *Indian J. Mar. Sci.*, 29: 48-51.
- Delvasto, P., A. Valverde, A. Ballester, J.M. Igual and J.A. Munoz *et al.*, 2006. Characterization of brushite as a re-crystallization product formed during bacterial solubilization of hydroxyapatite in batch cultures. *Soil Biol. Biochem.*, 38: 2645-2654.
- Donahue, R.L., R.W. Miller and J.C. Shickluna, 1990. An Introduction to Soils and Plant Growth. Prentice Hall of India Private Ltd., New Delhi, pp: 222.
- Garcia, L., A. Probanza, B. Ramos, J. Barriuso and F.J. Gutierrez Manero, 2004. Effects of inoculation with plant growth promoting rhizobacteria (PGPRs) and *Sinorhizobium fredii* on biological nitrogen fixation, nodulation and growth of *Glycine max* cv. Osumi. *Plant and Soil*, 267: 143-153.
- Girgis, M.G.Z., H.M.A. Khalil and M.S. Sharaf, 2008. *In vitro* evaluation of rock phosphate and potassium solubilizing potential of some *Bacillus* Strains. *Aust. J. Basic Applied Sci.*, 2: 68-81.

- Groudeva, V.I. and S.N. Groudev, 1987. Aluminosilicate biodegradation in the soil. Proceedings of the 9th International Symposium on Soil Biology and Conservation of the Biosphere, (ISSBCB'87), Akademiai Kiado, Budapest, pp: 621-628.
- Han, H., S. Supanjani and K.D. Lee, 2006. Effect of co-inoculation with phosphate and potassium solubilizing bacteria on mineral uptake and growth of pepper and cucumber. *Plant Soil Environ.*, 52: 130-136.
- Klopper, J.W., D.J. Hume, F.M. Scher, C. Singleton and B. Tipping *et al.*, 1988. Plant growth-promoting rhizobacteria on canola (rapeseed). *Plant Dis.*, 72: 42-46.
- Lebo, M.E., 1990. Phosphate uptake along a coastal plain estuary. *Limnol. Oceanogr.*, 35: 1279-1289.
- Lindsay, W.L., P.L.G. Vlek and S.H. Chien, 1989. Phosphate Minerals. In: Minerals in Soil Environment, Dixon, J.B. and S.B. Weed, (Ed.). Soil Science Society of America, Madison, USA., pp: 1089.
- Liu, W., X. Xu, X. Wu, Q. Yang, Y. Luo and P. Christie, 2006. Decomposition of silicate minerals by *Bacillus mucilaginosus* in liquid culture. *Environ. Geochem. Health*, 28: 133-140.
- Ouahmane, L., J. Thioulouse, M. Hafidi, Y. Prin and M. Ducouso *et al.*, 2007. Soil functional diversity and P solubilization from rock phosphate after inoculation with native or allochthonous arbuscular mycorrhizal fungi. *For. Ecol. Manage.*, 241: 200-208.
- Premono, E.M., A.M. Moawad and P.L.G. Vlek, 1996. Effect of phosphate solubilizing *Pseudomonas putida* on the growth of maize and its survival in the rhizosphere. *Indonesian J. Crop Sci.*, 11: 13-23.
- Rodriguez, H. and R. Fraga, 1999. Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnol. Adv.*, 17: 319-339.
- Rudresh, D.L., M.K. Shivaprakash and R.D. Prasad, 2005. Tricalcium phosphate solubilizing abilities of *Trichoderma* spp. in relation to P uptake and growth and yield parameters of chickpea (*Cicer arietinum* L.). *Can. J. Microbiol.*, 51: 217-222.
- Seshadri, S., S. Ignacimuthu and C. Lakshminarsimhan, 2002. Variations in heterotrophic and phosphate solubilizing bacteria from Chennai, southeast coast of India. *Indian J. Mar. Sci.*, 31: 69-72.
- Soltanpour, P.N. and S.M. Workman, 1979. Oil Testing Methods used at Colorado State University Soil Testing Laboratory for the Evaluation of Fertility, Salinity, Sodicity and Trace Element Toxicity. Colorado State University Experiment Station, Fort Collins, Colorado, USA., pp: 14.
- Styriakova, I., I. Styriak and T. Sasvari, 2004. Extraction of elements from sulphide and silicate concentrates by selected bacillus isolates. *Metalurgija*, 43: 293-297.
- Van, D. and P. Boers, 1994. Influence of internal loading on phosphorus concentration in shallow lakes before and after reduction of the external loading. *Hydrobiologia*, 275-76: 379-389.
- Venkateswaran, K. and R. Natarajan, 1983. Seasonal distribution of inorganic phosphate solubilizing bacteria and phosphatase producing bacteria in Porto Novo waters. *Indian J. Mar. Sci.*, 12: 213-217.
- Vikram, A., 2007. Efficacy of phosphate solubilizing bacteria isolated from vertisols on growth and yield parameters of sorghum. *Res. J. Microbiol.*, 2: 550-559.
- Vikram, A. and H. Hamzehzarghani, 2008. Effect of phosphate solubilizing bacteria on nodulation and growth parameters of greengram (*Vigna radiata* L. Wilczek). *Res. J. Microbiol.*, 3: 62-72.
- Zayed, G. and H. Abdel-Motaal, 2005a. Bio-active composts from rice straw enriched with rock phosphate and their effect on the phosphorus nutrition and microbial community in rhizosphere of cowpea. *Bioresour. Technol.*, 96: 929-935.
- Zayed, G. and H. Abdel-Motaal, 2005b. Bio-production of compost with low pH and high soluble phosphorus from sugar cane bagasse enriched with rock phosphate. *World J. Microbiol. Biotechnol.*, 21: 747-752.