



# Genome Sequence of *Coxiella*-Like Endosymbiont Strain CLE-RmD, a Bacterial Agent in the Cattle Tick (*Rhipicephalus microplus*) Deutsch Strain

Arunachalam Ramaiah,<sup>a</sup> Gregory A. Dasch<sup>a</sup>

<sup>a</sup>Rickettsial Zoonoses Branch, Centers for Disease Control and Prevention, Atlanta, Georgia, USA

**ABSTRACT** We report a partial genome sequence for the *Coxiella*-like endosymbiont strain CLE-RmD, assembled from metagenomics data obtained from the southern cattle tick (*Rhipicephalus microplus*) Deutsch strain.

Many of the model species for investigating host-bacterial interactions include hematophagous, plant sap-sucking arthropods, and food-storage pests (1–4). These diets lack essential amino acids, vitamins, and cofactors, which also cannot be synthesized by the host arthropods. The endosymbiotic bacteria of these arthropods are hypothesized to provide those nutrients (5). Bacterial endosymbionts in hard and soft ticks related to *Coxiella burnetii*, the etiologic agent of Q fever, are known as *Coxiella*-like endosymbionts (CLEs) (6–8). The nutritional dependence of *Amblyomma americanum* on its CLE has been experimentally demonstrated (9) and further supported bioinformatically for CLEs from several other hard tick species whose symbiont genomes have been sequenced, namely, *Rhipicephalus turanicus* (CRt) (10), *Rhipicephalus sanguineus* (CLE-RsOK) (11), and *A. americanum* (CLEAA/CLE-AaGA) (11, 12). For most other tick CLE symbionts, only a few sequences have been obtained (7).

Illumina HiSeq-1000 (accession numbers SRX1668952, SRX1668960, and SRX1668961) and 454GS FLX (accession numbers SRX019998 and SRX019999) reads were generated from low-Cot and Cot 696 genomic DNA extracted from a pooled collection of eggs from the f7 and f10 to f12 generations of the *R. microplus* Deutsch strain colony from Webb County, Texas (13–16), and submitted by the Centre for Comparative Genomics, Murdoch University, Australia (BioProject numbers PRJNA312025 and PRJNA46685, respectively). All of the Illumina runs and 454 reads were merged independently and mapped against 38 *Coxiella* genome sequences (NCBI/DDBJ/EMBL) and NCBI contigs containing CLE sequences previously assembled from Illumina/PacBio/454 reads of *R. microplus* using BWA v0.7.12 (17). All of the Illumina and 454 mapped CLE-RmD reads were extracted using SAMtools v1.3.1 (18) and independently assembled into scaffolds using the state-of-the-art assemblers SPAdes v3.10.1 (19), Velvet v1.2.10 (20), and CLC Genomics Workbench v9.5.2 (Qiagen). The CISA v1.3 tool (21) was used to integrate the three *de novo* assemblies with the NCBI *R. microplus* CLE contigs into a hybrid set of scaffolds, from which duplicate regions were excluded.

The partial genome of CLE-RmD (1,296,618 bp, 89 contigs with 229× coverage) was annotated with Prokka (22) and BLAST (23). The size of CLE-RmD (1.30 Mb, 31.7% G+C) is 65.0% of the size of the genome of the human pathogen *C. burnetii* (1.99 Mb, 42.7% G+C) but 81.9% of CRt (1.58 Mb, 38.2% G+C; we excluded the near-exactly duplicated 150,764-bp region from coordinates 590,925 to 741,688, which is probably an artifact) (10) and 116.2% that of the partial CLE-RsOK genome assembly (1.12 Mb, 38.0%) (11, 12). All three CLEs from *Rhipicephalus* (CRt, CLE-RsOK, and CLE-RmD) are much larger than the complete CLEAA and CLE-AaGA genomes from *Amblyomma* (657 kb, 34.6% G+C) (11, 12). The partial CLE-RmD genome contained 742 protein genes, 28 tRNAs, 3

Received 29 January 2018 Accepted 21 February 2018 Published 29 March 2018

**Citation** Ramaiah A, Dasch GA. 2018. Genome sequence of *Coxiella*-like endosymbiont strain CLE-RmD, a bacterial agent in the cattle tick (*Rhipicephalus microplus*) Deutsch strain. Genome Announc 6:e00003-18. <https://doi.org/10.1128/genomeA.00003-18>.

This is a work of the U.S. Government and is not subject to copyright protection in the United States. Foreign copyrights may apply. Address correspondence to Gregory A. Dasch, [ged4@cdc.gov](mailto:ged4@cdc.gov).

rRNAs, and 3 small RNAs (sRNAs) (transfer-messenger RNA [tmRNA], single recognition particle [SRP], and RNaseP). CLE-RmD genes typically had 80 to 90% similarity to homologues in CRt and CLE-RsOK but had at least 10% lower similarity to those of CLEAA, CLE-AaGA, and *C. burnetii*.

**Accession number(s).** The genome of *Coxiella* endosymbiont strain CLE-RmD of *R. microplus* was submitted as a whole-genome shotgun (WGS) project at GenBank/DDBJ/ENA under the accession number [DLUJ00000000](https://doi.org/10.1093/gbe/evv016). The WGS of the CLE-RmD version described in this paper is version DLUJ01000000.

## ACKNOWLEDGMENTS

A.R. was supported in part by an appointment in the Postdoctoral Bioinformatics Fellowship Program administered by the Association of Public Health Laboratories (APHL) and funded by the Centers for Disease Control and Prevention (CDC) as part of Cooperative Agreement 1U60OE000103 (CFDA number 93.322).

The findings and conclusions in this article are those of the authors and do not necessarily represent the views of the Centers for Disease Control and Prevention, U.S. Department of Health and Human Services.

## REFERENCES

- Buchner P. 1965. Endosymbiosis of animals with plant microorganisms. Interscience Publishers, New York, NY.
- Baumann P. 2005. Biology of bacteriocyte-associated endosymbionts of plant sap-sucking insects. *Annu Rev Microbiol* 59:155–189. <https://doi.org/10.1146/annurev.micro.59.030804.121041>.
- Moran NA, McCutcheon JP, Nakabachi A. 2008. Genomics and evolution of heritable bacterial symbionts. *Annu Rev Genet* 42:165–190. <https://doi.org/10.1146/annurev.genet.41.110306.130119>.
- Wernegreen JJ. 2012. Strategies of genomic integration within insect-bacterial mutualisms. *Biol Bull* 223:112–122. <https://doi.org/10.1086/BBLv223n1p112>.
- Rio RV, Attardo GM, Weiss BL. 2016. Grandeur alliances: symbiont metabolic integration and obligate arthropod hematophagy. *Trends Parasitol* 32:739–749. <https://doi.org/10.1016/j.pt.2016.05.002>.
- Noda H, Munderloh UG, Kurtz TJ. 1997. Endosymbionts of ticks and their relationship to *Wolbachia* spp. and tick-borne pathogens of humans and animals. *Appl Environ Microbiol* 63:3926–3932.
- Duron O, Noël V, McCoy KD, Bonazzi M, Sidi-Boumedine K, Morel O, Vavre F, Zenner L, Jourdain E, Durand P, Arnathau C, Renaud F, Trape JF, Biguezoton AS, Cremaschi J, Dietrich M, Léger E, Appelgren A, Dupraz M, Gómez-Díaz E, Diatta G, Dayo GK, Adakal H, Zoungrana S, Vial L, Chevillon C. 2015. The recent evolution of a maternally-inherited endosymbiont of ticks led to the emergence of the Q fever pathogen, *Coxiella burnetii*. *PLoS Pathog* 11:e1004892. <https://doi.org/10.1371/journal.ppat.1004892>.
- Reeves WK, Loftis AD, Sanders F, Spinks MD, Wills W, Denison AM, Dasch GA. 2006. *Borrelia*, *Coxiella*, and *Rickettsia* in *Carios capensis* (Acari: Argasidae) from a brown pelican (*Pelecanus occidentalis*) rookery in South Carolina, USA. *Exp Appl Acarol* 39:321–329. <https://doi.org/10.1007/s10493-006-9012-7>.
- Zhong J. 2012. *Coxiella*-like endosymbionts. *Adv Exp Med Biol* 984:365–379. [https://doi.org/10.1007/978-94-007-4315-1\\_18](https://doi.org/10.1007/978-94-007-4315-1_18).
- Gottlieb Y, Lalzar I, Klasson L. 2015. Distinctive genome reduction rates revealed by genomic analyses of two *Coxiella*-like endosymbionts in ticks. *Genome Biol Evol* 7:1779–1796. <https://doi.org/10.1093/gbe/evv108>.
- Ramaiah A, Williams-Newkirk AJ, Frace MA, Zambrano ML, Dasch GA. 2017. Evolution of the genomes of the *Coxiella*-like endosymbionts of ticks. *J Mississippi Acad Sci* 62:21. [http://msacad.org/wp-content/uploads/2017/02/Vol-62\\_Jan\\_1\\_Journal-of-MAS\\_abstract\\_Issue\\_2\\_1\\_17\\_b.pdf](http://msacad.org/wp-content/uploads/2017/02/Vol-62_Jan_1_Journal-of-MAS_abstract_Issue_2_1_17_b.pdf).
- Smith TA, Driscoll T, Gillespie JJ, Raghavan R. 2015. A *Coxiella*-like endosymbiont is a potential vitamin source for the Lone Star tick. *Genome Biol Evol* 7:831–838. <https://doi.org/10.1093/gbe/evv016>.
- Barrero RA, Guerrero FD, Black M, McCooke J, Chapman B, Schilkey F, Pérez de León AA, Miller RJ, Bruns S, Dobry J, Mikhaylenko G, Stormo K, Bell C, Tao Q, Bogden R, Moolhuijzen PM, Hunter A, Bellgard MI. 2017. Gene-enriched draft genome of the cattle tick *Rhipicephalus microplus*: assembly by the hybrid Pacific Biosciences/Illumina approach enabled analysis of the highly repetitive genome. *Int J Parasitol* 47:569–583. <https://doi.org/10.1016/j.ijpara.2017.03.007>.
- Guerrero FD, Moolhuijzen P, Peterson DG, Bidwell S, Caler E, Bellgard M, Nene VM, Djikeng A. 2010. Reassociation kinetics-based approach for partial genome sequencing of the cattle tick, *Rhipicephalus microplus*. *BMC Genomics* 11:374. <https://doi.org/10.1186/1471-2164-11-374>.
- Bellgard M, Moolhuijzen PM, Guerrero FD, Schibeci D, Rodriguez-Valle M, Peterson DG, Dowd SE, Barrero R, Hunter A, Miller RJ, Lew-Tabor AE. 2012. CattleTickBase: an integrated internet-based bioinformatics resource for *Rhipicephalus (Boophilus) microplus*. *Int J Parasitol* 42:161–169. <https://doi.org/10.1016/j.ijpara.2011.11.006>.
- McCooke JK, Guerrero FD, Barrero RA, Black M, Hunter A, Bell C, Schilkey F, Miller RJ, Bellgard MI. 2015. The mitochondrial genome of a Texas outbreak strain of the cattle tick, *Rhipicephalus (Boophilus) microplus*, derived from whole genome sequencing Pacific Biosciences and Illumina reads. *Gene* 571:135–141. <https://doi.org/10.1016/j.gene.2015.06.060>.
- Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* 26:589–595. <https://doi.org/10.1093/bioinformatics/btp698>.
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R; 1000 Genome Project Data Processing Subgroup. 2009. The sequence alignment/map (SAM) format and SAMtools. *Bioinformatics* 25:2078–2079. <https://doi.org/10.1093/bioinformatics/btp352>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <https://doi.org/10.1101/gr.074492.107>.
- Lin S-H, Liao Y-C. 2013. CISA: contig integrator for sequence assembly of bacterial genomes. *PLoS One* 8:e60843. <https://doi.org/10.1371/journal.pone.0060843>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403–410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2).